
TEST BANK FOR GENERAL CHEMISTRY 2 1 PRINCIPLES PATTERNS AND APPLICATIONS 2ND EDITION AVERILL ISBN 9781453336175





Chapter 2

Molecules, Ions, and Chemical Formulas

Section 2.1

True/False Questions

1. In an ionic bond, the atoms are held together by the electrostatic attraction There is a triple bond between the two oxygen atoms in a molecule of O2.
 False; Easy
 Methane (CH₄) is a covel between the positively charged nuclei of the bonded atoms and the negatively

- carbon and hydrogen atoms.

True; Easy

4. A Ca²⁺ ion has the same number of electrons as the noble gas Argon.

True; Easy

5. Ions that contain fewer electrons than protons have a net positive charge and are called anions.

False; Easy

Multiple Choice Questions

- 1. Which of the following is an elemental-polyatomic molecule?
 - a. Carbon tetrachloride
 - b. Nitrogen
 - c. Phosphorous
 - d. Sodium chloride
 - e. Chlorine
 - c; Easy
- 2. How many electrons are shared between two atoms joined by a double bond?
 - a. One
 - b. Two
 - c. Four
 - d. Six
 - e. Eight
 - c; Easy
- 3. Which of the following is an organic compound?
 - FeO a.
 - NaHCO₃ b.
 - CH_4 c.
 - d. SF_6

	e. H ₂ CO ₃ c; Easy
4.	Which of the following compounds has only covalent bonding? a. H_2O b. $NaOH$ c. $NaCl$ d. $Al_2(SO_3)_3$ e. Fe_2O_3
5.	Which of the following molecules has a tetrahedral geometry? a. H_2O b. CO_2 c. CH_4 d. NO_2 e. NH_3 c; Easy Which molecular representation method uses wedges and dashes to sketch the
6.	Which molecular representation method uses wedges and dashes to sketch the structure of a molecule in three dimensions? a. Ball-and-stick model b. Perspective drawing c. Space filling model d. Condensed structural formula e. Molecular formula b; Easy
7.	If two charged particles, Q_1 and Q_2 , are separated by the distance r , the electrostatic energy between the particles is ∞ a. $Q_1 \times Q_2 + r$ b. $Q_1 \times Q_2 \times r$ c. $\frac{r^2}{Q_1 \times Q_2}$ d. $\frac{Q_1Q_2}{r}$ e. $\frac{r}{Q_1 \times Q_2}$
8.	The most common monoatomic ion formed by magnesium is a. Mg^- b. Mg^{2-} c. Mg^{3+} d. Mg^{2+} e. Mg^{3-}

© 2021 FlatWorld

9. What is the total number of electrons present in an Mg^{2+} ion?

a. 8
b. 10
c. 12
d. 14
e. 16
b; Easy

Essay Questions

1. Briefly define the term chemical bond and then describe the types of chemical bonds.

Atoms form chemical compounds when the attractive electrostatic interactions between them are stronger than the repulsive interactions. Collectively, we refer to the attractive interactions between atoms as chemical bonds. Chemical bonds are generally divided into two fundamentally different kinds: ionic and covalent. In reality, however, the bonds in most substances are neither purely ionic nor purely covalent, but they are closer to one of these extremes. Ionic compounds consist of positively and negatively charged ions held together by strong electrostatic forces, whereas covalent compounds generally consist of molecules, which are groups of atoms in which one or more pairs of electrons are shared between bonded atoms. **Easy**

2. Differentiate between organic and inorganic compounds.

Covalent compounds that contain predominantly carbon and hydrogen are called organic compounds. Compounds that consist primarily of elements other than carbon and hydrogen are called inorganic compounds; they include both covalent and ionic compounds.

Easy

Fill in the Blank

1.	In a(n), the atoms are held together by the electrostatic attraction between
	the positively charged nuclei of the bonded atoms and the negatively charged
	electrons they share.
	covalent bond; Easy
	76,
2.	Each covalent compound is represented by a(n), which gives the atomic
2	symbol for each component element, in a prescribed order, accompanied by a
	subscript indicating the number of atoms of that element in the molecule.
	molecular formula; Easy
3.	Ions that contain more electrons than protons have a net negative charge and are
	called
	anions; Easy
4.	Cl^- is $a(n)$ ion.
	monoatomic; Moderate

Chlorine with atomic number 17 would most likely an electron, during ionic bonding, to attain the noble gas configuration of Argon. gain; Easy
<u>ction 2.2</u>
rue/False Questions
The formula unit of a compound is the relative numbers of atoms of the elements in a compound, reduced to the smallest whole numbers. False; Easy
False; Easy Sodium oxide (Na ₂ O) is a binary ionic compound. True; Easy
The thiocyanate ion (SCN ⁻) is an example of a polyatomic anion. True; Easy
The dichromate ion (Cr ₂ O ₇ ⁻) is an example of a polyatomic cation. False; Easy
The ion pair Na ⁺ and SO ₄ ²⁻ forms the compound Na ₄ SO. False; Easy
ultiple Choice Questions
The compound formed from Fe ³⁺ and O ²⁻ is a. FeO b. Fe ₂ O c. FeO ₂ d. FeO ₃ e. Fe ₂ O ₃ e; Easy
Which of the following is the compound formed from the ions Pb^{2+} and S^{2-} ? a. PbS b. Pb_2S c. PbS_2 d. $2Pb_2S$
e. Pb ₂ S ₂ a; Easy
The empirical formula for the compound formed from an aluminum ion and a sulfate ion is a. AlSO ₄ b. Al(SO ₄) ₂ c. Al ₂ SO ₄ d. (AlSO ₄) ₂ e. Al ₂ (SO ₄) ₃

	e; Easy
4.	Which of the following is the compound formed from the ions Cu^+ and O^{2-} ? a. CuO b. Cu_2O c. CuO_2 d. CuO_3 e. Cu_2O_3
5.	Which of the following is the formula of the polyatomic ion peroxide? a. $O_2^{2^-}$ b. CN^- c. $HPO_4^{2^-}$ d. $HPO_4^{2^-}$ e. $HPO_3^{2^-}$ a; Easy Which of the following is the compound formed from the ions NH_4^+ and NO_3^- ?
6.	Which of the following is the compound formed from the ions NH ₄ ⁺ and NO ₃ ⁻ ? a. N ₂ O ₂ H ₂ b. HNO ₃ c. NO ₄ NH ₃ d. NH ₃ NO ₄ e. NH ₄ NO ₃ e; Easy say Questions What are polyatomic ions? Give an example of a polyatomic anion and a
Es	say Questions
	What are polyatomic ions? Give an example of a polyatomic anion and a polyatomic cation. Polyatomic ions are groups of atoms that bear a net electrical charge, although the atoms in a polyatomic ion are held together by the same covalent bonds that hold atoms together in molecules. Just as there are many more kinds of molecules than simple elements, there are many more kinds of polyatomic ions than monatomic ions. An example of a polyatomic cation is the ammonium ion with formula NH ₄ An example of a polyatomic anion is the sulfate ion with formula SO ₄ ²⁻ . Easy
Fil	l in the Blank
1.	The of a compound is the relative number of atoms of the elements in a compound, reduced to the smallest whole numbers. empirical formula; Easy
2.	An ionic compound that contains only two elements, one present as a cation and one as an anion, is called $a(n)$ binary ionic compound; Easy

3.	HSO ₄ ⁻ is an example of a polyatomic anion; Easy						
4.	A dichromate ion, $Cr_2O_7^{2-}$, will react with potassium ions to form potassium dichromate. two; Easy						
5.	are compounds that contain specific ratios of loosely bound water molecules. Hydrates; Easy						
Sec	ue/False Questions						
Tr	ue/False Questions						
1.							
2.	Two chlorine ions react with one chromium (II) ion to form chromic chloride. False; Easy						
3.	NaClO ₄ is referred to as sodium hypochlorite. False; Easy NaC ₂ H ₃ O ₂ is called sodium acetate.						
4.	NaC ₂ H ₃ O ₂ is called sodium acetate. True; Easy						
Μι	ultiple Choice Questions						
1.	The common name for the compound Fe ₂ (SO ₄) ₃ is a. iron sulfate b. ferric sulfate c. iron-two sulfate d. ferrous sulfate e. iron-three sulfate b; Easy						
2	What is the systematic name of NH ₄ ClO ₄ ?						
	a. Chlorine nitride						
0	b. Ammonium chlorate						
	c. Ammonium chloride						
	d. Chlorammonium hydride						
	e. Ammonium perchlorate e; Easy						
2	Will all the state of the Co						
3.	•						
	a. Ferric sulfide						
	b. Iron(I) sulfidec. Iron (II) sulfide						
	v. Hon (ii) builde						

	d. Ferrous sulfide
	e. Iron (III) sulfide
	c; Easy
4.	The formula for sodium bicarbonate is
	a. NaCO ₂
	b. Na_2CO_3
	d. NaHCO ₃
	e. $Na_2(CO_3)_2$
	d; Easy
5.	c. Na_2HCO_3 d. $NaHCO_3$ e. $Na_2(CO_3)_2$ d; Easy The ion $Cr_2O_7^{2-}$ is called a. per chromate b. chromite c. chromium oxate d. chromate e. dichromate e. dichromate e; Easy The formula for the sulfite ion is a. SO_3^{2-} b. SO_4^{2-} c. SO_2^{-} d. SO_2^{2-} e. SO_2^{4-} a; Easy The systematic name of $Mg(HCO_3)_2$ is
	a. per chromate
	b. chromite
	c. chromium oxate
	d. chromate
	e. dichromate
	e; Easy
6.	The formula for the sulfite ion is
	a. SO_3^{2-}
	b. SO_4^{2-}
	c. $SO_2^{\frac{1}{2}}$
	d. SO_{3}^{2}
	e. SO_2^{4-}
	or Fooy
	a; Easy
7.	The systematic name of Mg(HCO ₃) ₂ is
	a. magnesiam careonate
	b. magnesium bicarbonate
	c. magnesium formate
	d. magnesium acetate
	e. manganese (II) bicarbonate
	b; Easy
8 -	The formula for aluminum sulfite is
3.	a. AlSO ₃
6	b. $Al_2(SO_3)_2$
-	c. $Al_2(SO_4)_3$
	d. $Al_3(SO_3)_2$
	e. $Al_2(SO_3)_3$
	e; Easy

Essay Questions

1. Explain the different conventions involved in naming a polyatomic anion.

Polyatomic anions that contain a single metal or nonmetal atom plus one or more oxygen atoms are called oxoanions. In cases where only two oxoanions are known for an element, the name of the oxoanion with more oxygen atoms ends in -ate, and the name of the oxoanion with fewer oxygen atoms ends in -ite. For example, NO₃⁻ is nitrate and NO₂⁻ is nitrite. The halogens and some of the transition metals form more extensive series of oxoanions with as many as four members. In the names of these oxoanions, the prefix per- is used to identify the oxoanion with the most oxygen (so that ClO₄⁻ is perchlorate and ClO₃⁻ is chlorate), and the prefix hypo- is used to identify the anion with the fewest oxygen (ClO₂⁻ is chlorite and oissinneu.ca7 ClO⁻ is hypochlorite).

Moderate

Fill in the Blank

	• • • • • • • • • • • • • • • • • • • •
1.	Polyatomic anions that contain a single metal or nonmetal atom plus one or more oxygen atoms are called oxyanions; Easy
2.	The common or old name for iron (II) sulfide is
	ferrous sulfide; Easy
3.	(NH ₄) ₂ SO ₄ is called in the systematic nomenclature.
	ammonium sulfate; Easy
4.	The formula for calcium formate is
	Ca(HCO ₂) ₂ ; Easy

Section 2.4

True/False Questions

True; Easy

- 1. The formula of nitrogen dioxide is N_2O_2 . False; Easy
- 2. The systematic name for OF_2 is oxygen difluoride.
- 3. Cyclohexane is a saturated hydrocarbon.
- True; Easy
- 4. Cyclopropane has four carbon atoms in its structure. False; Easy
- 5. C₆H₅CH₂OH is the condensed molecular formula of phenol. False; Easy

Multiple Choice Questions

1. What is the name of RbS?

	a. Sulfo rhenium
	b. Rhodium sulfide
	c. Sulfo ruthenium
	d. Ruthenium sulfide
	e. Rubidium sulfide
	e; Easy
2.	What is the name of Tl ₂ Se?
	a. Titanium (II) sulfate
	b. Titanium (I) sulfite
	c. Terillium (II) selenide
	d. Thallium (I) selenide
	e. Tungsten (II) sulfide
	d; Moderate
3.	Which of the following is the formula of niobium (IV) oxide?
	a. Nb ₄ O
	b. Ni ₄ O
	c. NO_2
	d. N_2O_2
	e. NbO ₂
	d. Thallium (I) selenide e. Tungsten (II) sulfide d; Moderate Which of the following is the formula of niobium (IV) oxide? a. Nb ₄ O b. Ni ₄ O c. NO ₂ d. N ₂ O ₂ e. NbO ₂ e; Easy The formula of copper (I) oxide is
4.	The formula of copper (I) oxide is
	a. CuO
	b. Cu ₂ O
	c. Co_2O_2
	d. Cu ₃ O ₂
	e. CoO
	The formula of copper (I) oxide is
5.	Which of the following is the condensed structural formula for isobutane?
	a. CH ₃ CH ₂ CH ₂ CH ₃
	b. CH ₃ CH ₂ CH ₂
	c. CH ₃ CH ₂ CH ₂ CH ₃
	d. (CH ₃) ₃ CH ₂
	e. (CH ₃) ₂ CHCH ₃
-2	e; Easy
6.	Which of the following is the simplest alkene?
	a. C_2H_6
	b. C_6H_6
	c. C ₂ H ₄
	d. C_2H_2
	e. C_6H_{12}
	c; Moderate
7.	Which of the following is the structure of cyclobutane?
•	a.







c.



d.






d; Moderate

- 8. What is the name of $CH_3C \equiv CCH_3$?
 - 2-butane a.
 - 2-butene b.
 - 2-pentene c.
 - 2-butyne
 - 2-cyclobutane e.
 - d; Easy
- than phiterera apnathannonipissinneut. Cat 9. Which of the following is the structure of toluene?





c.



Essay Questions

1. Briefly describe the four major classes of hydrocarbons. What are saturated and unsaturated hydrocarbons?

The four major classes of hydrocarbons are the alkanes, which contain only carbon—hydrogen and carbon—carbon single bonds; the alkenes, which contain at least one carbon—carbon double bond; the alkynes, which contain at least one carbon—carbon triple bond; and the aromatic hydrocarbons, which usually contain rings of six carbon atoms that can be drawn with alternating single and double bonds. Alkanes are also called saturated hydrocarbons, whereas hydrocarbons that contain multiple bonds (alkenes, alkynes, and aromatics) are unsaturated. **Easy**

2. What are cyclic hydrocarbons? Provide examples of cyclic hydrocarbons, and then explain how to draw cyclic hydrocarbons.

In a cyclic hydrocarbon, the ends of a hydrocarbon chain are connected to form a ring of covalently bonded carbon atoms. Cyclic hydrocarbons are named by attaching the prefix cyclo- to the name of the alkane, the alkene, or the alkyne. The simplest cyclic alkanes are cyclopropane (C_3H_6) a flammable gas that is also a powerful anesthetic, and cyclobutane (C_4H_8). The most common way to draw the structures of cyclic alkanes is to sketch a polygon with the same number of vertices as there are carbon atoms in the ring; each vertex represents a CH_2 unit. **Easy**

Fill in the Blank

1.0		are the	simples	t class of	organi	c compound	ls, consisting	gentirely (of carbon
0	and hy	drogen							
		Ţ	_						

Hydrocarbons; Easy

2.	Unsaturated hydrocarbons are those which contain multiple bonds such as alkenes
	and
	alkynes; Easy

3.	CH ₃ CH ₂ CH ₂ CH ₂ CH ₃ is called
	n-pentane; Easy

4.	The molecular formula for cyclobutane is C4H8; Easy					
5.	The molecular formula of ethanol is CH ₃ CH ₂ OH; Easy					
Sec	etion 2.5					
Tr	ue/False Questions					
1.	Sulfuric acid is an example of an oxoacid. True; Easy HClO is the formula of perchloric acid. False; Easy Chlorous acid is the acid formed by adding one proton to the chlorite ion (ClO ₂ ⁻).					
2.	HClO is the formula of perchloric acid. False; Easy					
3.	Chlorous acid is the acid formed by adding one proton to the chlorite ion (ClO_2^-) . True; Easy					
4.	Calcium oxide when dissolved in water results in a basic solution. True; Easy					
5.	C ₂ H ₅ CHO is an example of carboxylic acids. False; Easy altiple Choice Questions					
Μι	ultiple Choice Questions					
1.	Which of the following is an oxoacid? a. Hydrochloric acid b. Hypochlorous acid c. Hydrocyanic acid d. Hydrosulphuric acid e. Hydroiodic acid b; Easy					
2.	H ₂ SO ₃ is the formula of a. persulfuric acid b. hyposulfurous acid					
60	c. sulfurous acid d. hydrosulfuric acid e. sulfuric acid c; Easy					
3.	Chloric acid is represented by the formula a. HClO b. HClO ₂ c. HClO ₃ d. HClO ₄ e. HCl					

	c; Easy
4.	Which of the following acids is a carboxylic acid? a. HClO b. HCOOH c. HCN d. HBr e. H ₂ S b; Easy
5.	HNO ₃ is the name of a. hydronitric acid b. hyponitrous acid c. nitrous acid d. pernitric acid e. nitric acid e; Easy Which of the following when added to water would result in a strongly basic
6.	Which of the following when added to water would result in a strongly basic solution? a. HClO b. NaCl c. HCN d. MgO e. CH ₃ OH d; Easy Which of the following forms a neutral solution when added to water?
7.	Which of the following forms a neutral solution when added to water? a. NaOH b. CH_3OH c. HCN d. $Mg(OH)_2$ e. H_2S b; Easy
8.	Which of the following contains a carbonyl group? a. HClO b. Ca(OH) ₂ c. HCN d. CH ₃ COOH e. H ₂ SO ₄ d; Easy

Essay Questions

1. Explain the naming of acids in which the proton is attached to an element other than oxygen, and then provide examples of those types of acids.

In acids where the H^+ ion is attached to some element (other than oxygen), the name of the acid begins with hydro- and ends in -ic, with the root of the name of

the other element or ion in between. The name of the anion derived from this kind of acid always ends in -ide. Thus hydrogen chloride (HCl) gas dissolves in water to form hydrochloric acid (which contains H⁺ and Cl⁻ ions), hydrogen cyanide (HCN) gas forms hydrocyanic acid (which contains H⁺ and CN⁻ ions), and so forth. Examples of this kind of acid are commonly encountered and very important. For instance, your stomach contains a dilute solution of hydrochloric acid to help digest food.

Easy

2. How do amines form, and what is their physiological importance?

Replacing a hydrogen atom of NH₃ with an alkyl group results in an amine (RNH₂), which is also a base. Amines have pungent odors—for example, methylamine (CH₃NH₂) is one of the compounds responsible for the foul odor associated with spoiled fish. The physiological importance of amines is suggested in the word vitamin, which is derived from the phrase vital amines. The word was coined to describe dietary substances that were effective at preventing scurvy, rickets, and other diseases because these substances were assumed to be amines. Subsequently, some vitamins have indeed been confirmed to be amines. **Easy**

Fil	l in the Blank
1.	A base is a substance which produces one or more ions and a cation when dissolved in aqueous solutions. hydroxide; Easy
2.	The common name for the acid with the condensed molecular formula HCO ₂ H is acid. formic; Easy
3.	When a proton is added to the sulfite ion (SO ₃ ²⁻) acid is formed. sulfurous acid; Easy
4.	Replacing a hydrogen atom of NH ₃ with an alkyl group results in a(n) amine; Easy
Se	ction 2.6
Te	ua/Falsa Quastions


1. In a refining tower, the less volatile fractions in petroleum condense near the top of the tower where it is cooler.

False; Easy

2. During cracking, less volatile, lower-value fractions are converted to more volatile, higher-value mixtures.

True; Easy

3. The higher the octane rating, the higher quality the fuel.



- 87 b.
- 58 c.
- 78 d.
- 190 e.
- d; Moderate
- 6. In the _____ process, water at high temperature (160°C) and high pressure is pumped underground to melt the sulfur, and compressed air is used to force the no hipisinne lica? liquid sulfur-water mixture to the surface.
 - a. contact
 - lead-chamber b.
 - c. Frasch
 - Haber's d.
 - fuming
 - c; Easy

Essay Questions

1. What is the octane scale, and how are fuels assigned octane ratings?

The octane scale was established in 1927 using a standard test engine and two pure compounds: *n*-heptane and isooctane (2, 2, 4-trimethylpentane). *n*-Heptane, which causes a great deal of knocking on combustion, was assigned an octane rating of 0, whereas isooctane, a very smooth-burning fuel, was assigned an octane rating of 100. Chemists assign octane ratings to different blends of gasoline by burning a sample of each in a test engine and comparing the observed knocking with the amount of knocking caused by specific mixtures of *n*-heptane and isooctane. For example, the octane rating of a blend of 89% isooctane and 11% nheptane is simply the average of the octane ratings of the components weighted by the relative amounts of each in the blend. Converting percentages to decimals, we obtain the octane rating of the mixture:

0.89(100) + 0.11(0) = 89

A gasoline that performs at the same level as a blend of 89% isooctane and 11% *n*-heptane is assigned an octane rating of 89.

Moderate

2. Explain the production of sulfuric acid using elemental sulfur as the starting material.

If sulfur is the starting material, the first step in the production of sulfuric acid is the combustion of sulfur with oxygen to produce SO₂. Next, SO₂ is converted to SO₃ by the contact process, in which SO₂ and O₂ react in the presence of V₂O₅ to achieve about 97% conversion to SO₃. The SO₃ can then be treated with a small amount of water to produce sulfuric acid. Usually, however, the SO₃ is absorbed in concentrated sulfuric acid to produce oleum, a more potent form called fuming sulfuric acid. Because of its high SO₃ content (approximately 99% by mass), oleum is cheaper to ship than concentrated sulfuric acid. At the point of use, the oleum is diluted with water to give concentrated sulfuric acid (very carefully because dilution generates enormous amounts of heat). Because SO₂ is a pollutant,

the small amounts of unconverted SO_2 are recovered and recycled to minimize the amount released into the air. Easy

Fill in the Blank

1.	Fuels having a(n) octane rating combust prematurely causing engine knocking or pinging. low; Easy
2.	A mixture of 77% isooctane and 23% <i>n</i> -heptane has an octane rating of 77 ; Easy
3.	In the process, sulfur was burned in a large room lined with lead, and the resulting fumes were absorbed in water. lead-chamber; Easy
Cles	77; Easy In the process, sulfur was burned in a large room lined with lead, and the resulting furnes were absorbed in water. lead-chamber; Easy